# Historical DNA from museum type specimens clarifies diversity of Asian leaf turtles (*Cyclemys*)

BRYAN L. STUART<sup>1\*</sup> and UWE FRITZ<sup>2</sup>

<sup>1</sup>The Field Museum, Department of Zoology, Division of Amphibians & Reptiles, 1400 S. Lake Shore Drive, Chicago, IL 60605-2496, USA <sup>2</sup>Museum of Zoology, Natural History State Collections Dresden, A. B. Meyer Building, Königsbrücker Landstr. 159, D-01109 Dresden, Germany

Received 22 February 2007; accepted for publication 9 July 2007

Species boundaries in Asian leaf turtles of the genus *Cyclemys* are difficult to define on the basis of morphology, primarily because many populations exhibit considerable ontogenetic variation in shell and head coloration. Two recent molecular phylogenetic hypotheses of *Cyclemys* species relationships, based largely on market and pet-trade samples of uncertain provenance, were highly incongruent. We used historical DNA methods to sequence fragments of the mitochondrial cytochrome *b* gene from eight type specimens of *Cyclemys* (including one collected by Alfred Russel Wallace), and phylogenetic hypothesis supports the recognition of four named species (*Cyclemys atripons*, *Cyclemys dentata*, *Cyclemys oldhamii*, and *C. pulchristriata*), as well as a fifth species of unknown geographical provenance obtained from the Hong Kong pet trade. The type sequences show that previous molecular phylogenetic studies were hampered by misidentifications, supporting the notion that *Cyclemys* of unknown provenance are not reliably identified to species solely on the basis of morphology. © 2008 The Linnean Society of London, *Biological Journal of the Linnean Society*, 2008, **94**, 131–141.

ADDITIONAL KEYWORDS: Geoemydidae – historical DNA – museum specimens – species delimitation.

## INTRODUCTION

Phylogenetic analyses of DNA sequence data are now regularly incorporated into taxonomic studies, and are especially useful in cases when species boundaries and evolutionary relationships are obscured by a high degree of morphological homoplasy. The most accurate means of matching DNA sequences to Linnaean names entail obtaining DNA sequence data from the type specimens on which the original descriptions of species were based (Tautz et al., 2003; Austin & Melville, 2006). However, most existing type specimens were collected prior to the molecular revolution in systematic biology, and so lack associated tissue samples that were specially preserved for genetic analysis. Methods to retrieve DNA from degraded samples remain challenging (Handt et al., 1994; Wayne, Leonard & Cooper, 1999; Cooper & Poinar, 2000; Hofreiter et al., 2001), but have transformed traditional museum specimens

Asian leaf turtles of the genus *Cyclemys* provide an excellent example of a taxonomic group in which species boundaries and their evolutionary relationships are obscured by a high degree of morphological homoplasy. Species of *Cyclemys* are primarily diagnosed by shell and head coloration, but many

into valuable sources of genetic material for systematic studies on rare and extinct populations and species (e.g. Roy *et al.*, 1994; Sorenson *et al.*, 1999; Austin & Arnold, 2001; Payne & Sorenson, 2002; Austin, Arnold & Jones, 2004; Kearney & Stuart, 2004; Parham *et al.*, 2004; Wisely, Maldonado, & Fleischer, 2004; Fuchs, Fjeldså & Pasquet, 2005; Asher & Hofreiter, 2006; Fleischer *et al.*, 2006; Kohlsdorf & Wagner, 2006). Relatively few systematic studies have attempted to obtain DNA from old type specimens (e.g. Krings *et al.*, 1997; Amato *et al.*, 1999; Hughey, Silva & Hommersand, 2001; Austin, Arnold & Bour, 2002; Austin, Arnold & Bour, 2003; Hassanin & Ropiquet, 2004; Johnson, Watson & Mindell, 2005; Stuart, Inger & Voris, 2006a).

<sup>\*</sup>Corresponding author. E-mail: bstuart@berkeley.edu

populations exhibit considerable ontogenetic variation in these characters (Fritz, Gaulke & Lehr, 1997; Fritz & Ziegler, 1999). Furthermore, some species of *Cyclemys* were originally described from juvenile specimens, and others were described from adult specimens (Fritz *et al.*, 1997; Iverson & McCord, 1997), making comparisons difficult. As a result, species boundaries in *Cyclemys* are contentious and notoriously difficult to define, and the genus has a complicated taxonomic history.

The genus Cyclemys is widely distributed from north-eastern India to the Philippines (Fritz et al., 1997; Iverson & McCord, 1997; Fritz & Ziegler, 1999). Historically, only one species, Cyclemys dentata (Gray, 1831), or sometimes two species, C. dentata and Cyclemys tcheponensis (Bourret, 1939) were thought to encompass this vast range. In 1997, a taxonomic revision (Fritz et al., 1997) and a new species description (Iverson & McCord, 1997) brought the number of recognized species to five (Artner, 1998; Fritz & Ziegler, 1999): Cyclemys atripons Iverson & McCord, 1997, with a type locality of Krat, southeastern Thailand; Cyclemys dentata (Gray, 1831), with a type locality of Java, Indonesia; Cyclemys oldhamii Gray, 1863, with a type locality of Mergui, Myanmar; Cyclemys pulchristriata Fritz, Gaulke & Lehr, 1997, with a type locality of Phuc-Son in central Vietnam; and Cyclemys tcheponensis (Bourret, 1939), with a type locality of Haute Sé-Bang-Hien on the border of southern Laos and central Vietnam. In the most recent taxonomic treatment, Guicking et al. (2002) resurrected Cyclemys dhor shanensis Annandale, 1918 out of synonymy with C. oldhamii, elevated shanensis to full species status, referred populations from Myanmar and Thailand to the subspecies Cyclemys shanensis shanensis Annandale, 1918, and referred C. tcheponensis to the subspecies Cyclemys shanensis tcheponensis (Bourret, 1939). Four additional described species of *Cyclemys* are currently considered to be junior synonyms of C. dentata or C. s. tcheponensis (reviewed in Fritz et al., 1997). Specifically, Cyclemys orbiculata Bell, 1834 and Cyclemys bellii Gray, 1863, with uncertain type localities, and Cyclemys ovata Gray, 1863, with a type locality of Sarawak are treated as junior synonyms of C. dentata, and Cyclemys tiannanensis Kou, 1989, with a type locality of Yunnan, China, is treated as a junior synonym of C. s. tcheponensis.

Despite the uncertainties in the number of taxa (species and subspecies), *Cyclemys* are readily classified into two distinct morphological groups on the basis of plastron (ventral part of the shell) coloration: *C. oldhamii*, *C. s. shanensis*, and *C. s. tcheponensis* have mostly dark brown or black plastra, whereas *C. atripons*, *C. dentata*, and *C. pulchristriata* have mostly yellow plastra (Fritz *et al.*, 1997; Guicking et al., 2002). Within the dark-plastron group, only adults of C. oldhamii and C. s. shanensis are reliably distinguished from C. s. tcheponensis, as C. oldhamii and C. s. shanensis lose head and neck stripes during ontogeny, but C. s. tcheponensis does not (Fritz et al., 1997; Guicking et al., 2002). No morphological characters have been reported that can reliably distinguish C. oldhamii from C. s. shanensis. Within the yellow plastron group, C. atripons and C. pulchristriata have immaculate throats, but C. dentata has a striped throat (Fritz et al., 1997; Iverson & McCord, 1997). No morphological characters have been reported that can reliably distinguish C. atripons from C. pulchristriata. Fritz et al. (2001) proposed that the number of ventral neck stripes differed between C. atripons and C. pulchristriata, but the range of neck stripes reported by Stuart & Platt (2004) in just two turtles collected together in southwestern Cambodia (near to the type locality of C. atri*pons*) exceeded the range reported for both species by Fritz et al. (2001). Because of their morphological similarity, some authors have suggested that C. atripons and C. pulchristriata may be conspecific (Fritz & Obst, 1999; Iverson in Guicking et al., 2002).

Defining species boundaries in Cyclemys is more than academic. Asian turtles are overexploited for food, for traditional medicine, and for pets (van Dijk, Stuart & Rhodin, 2000), and now more than half of the Asian species are listed as endangered or critically endangered (IUCN, 2004). The conservation status of Cyclemys has been evaluated as a single, widespread taxon, C. dentata, owing to the uncertainty in the limits and geographical distributions of the other named species (IUCN/SSC Tortoise and Freshwater Turtle Specialist Group and Asian Turtle Trade Working Group, 2000). However, the genus Cyclemys contains multiple, highly divergent mitochondrial DNA lineages (Guicking et al., 2002; Spinks et al., 2004), each of which implicitly have smaller geographical ranges that are more vulnerable to extinction than a single evolutionary lineage of Cyclemys. A more realistic assessment of species boundaries in Cyclemys will allow for a more accurate, and probably direr, assessment of the conservation status of these turtles.

An independent molecular dataset is highly desirable given the uncertainties in defining species of *Cyclemys* solely on the basis of morphology. Two previous hypotheses of phylogenetic relationships of *Cyclemys* species have been proposed, both based on the mitochondrial cytochrome b (cyt b) gene. Guicking *et al.* (2002) found samples of *C. oldhamii*, *C. atripons*, and *C. pulchristriata* to be paraphyletic, and interpreted a lineage morphologically resembling *C. oldhamii*, and a lineage morphologically resembling *C. atripons* and *C. pulchristriata*, to be cryptic, undescribed species. Spinks et al. (2004) included one sample of each of three species, and found *C. atripons* to be the sister taxon to a clade containing C. dentata and C. tcheponensis, with C. dentata and C. tcheponensis separated by a very short branch. This hypothesis implies that species with yellow plastra do not represent a monophyletic group. However, these studies relied entirely (Spinks et al., 2004) or mostly (Guicking et al., 2002) on pet-trade and market samples of uncertain provenance. For example, Guicking et al. (2002) assigned the names C. atripons and C. pulchristriata to samples depending on whether they were obtained from Cambodian markets (C. atripons) or Vietnamese markets (C. pulchristriata), despite the fact that turtles collected in Cambodia are mass exported to Vietnam (Stuart et al., 2000). Pet-trade and market samples do have utility in phylogenetic studies, but with serious limitations (Parham et al., 2001, 2004; Stuart & Parham, 2004). The uncertainties of identifying Cyclemys species based on morphology makes the limitations of using market and pet-trade samples even more severe, and the hypotheses of species boundaries and relationships of *Cyclemys* proposed in both studies need to be interpreted with caution. Clearly, DNA sequences from type specimens of *Cyclemys* would help resolve taxonomic uncertainty.

In an effort to elucidate species boundaries in Cyclemys, we used historical (or 'ancient') DNA methods to sequence fragments of the mitochondrial cyt b gene from museum type specimens, and phylogenetically placed these type sequences into the context of published cyt b variation. Although mitochondrial gene trees may not always represent species trees owing to introgression or lineage sorting of ancestral polymorphism (Moore, 1995; Maddison, 1997), an accurate matching of Linnaean names to major mitochondrial lineages provides an evidencebased hypothesis of species boundaries that is superior to the current state of taxonomic upheaval in Cyclemys. Furthermore, the hypothesis that major mitochondrial DNA lineages of Cyclemys represent species is more likely to overestimate diversity than to underestimate diversity, a source of error that we are willing to assume in this group of threatened turtles.

# MATERIAL AND METHODS

## EXTRACTION, AMPLIFICATION, AND SEQUENCING OF HISTORICAL DNA FROM MUSEUM SPECIMENS

Small pieces of bone, dried muscle, or connective tissue were removed from the inside of seven dried (stuffed or shell) type specimens of *C. atripons*, *C. bellii*, *C. oldhamii*, *C. orbiculata*, and *C. ovata*, and

a small piece of skin, muscle, or femur bone was removed from the leg of seven alcoholic type specimens of C. dentata and C. pulchristriata (Table 1). DNA was extracted following the method described by Kearney & Stuart (2004). Three fragments of 144, 179, and 168 nucleotide basepairs (bp; after primer sequences were trimmed) of mitochondrial DNA encoding part of the cyt b gene were amplified by polymerase chain reaction (PCR; 94 °C 45 s, 48 °C 30 s, 72 °C 1 min) for 40 cycles using the primer pairs L-350cvtb/H-500cvtb, L-430cvtb/H-615cvtb, and L-560cytb/H-730cytb, respectively (Table 2). Primers were designed so that the three fragments overlapped by 64 and 51 bp after primer sequences were trimmed. This measure ensured that fragments of contaminant DNA were not concatenated into chimeric sequences that might be erroneously judged to be authentic (Olson & Hassanin, 2003). The 25-µL PCR reactions utilized 4 µL of bovine serum albumin (BSA; New England BioLabs), to prevent PCR inhibitors, a relatively large volume (4 µL) of DNA template to overcome low extraction vield, the highquality Taq polymerase AmpliTaq Gold (Roche), and extra cycles (40 total) of the PCR reaction. All PCR reactions contained a positive control (described below) and a negative control (all reagents except the DNA template). PCR products were electrophoresed in a 1% low-melt agarose TALE gel stained with ethidium bromide and visualized under ultraviolet light. The bands containing DNA were excised, and agarose was digested from the bands using GELase (Epicentre Technologies). PCR products were sequenced in both directions by direct double-strand cycle sequencing using Big Dye version 3 (Perkin Elmer). The amplifying primers were used in the sequencing reactions. Cycle-sequencing products were precipitated with ethanol, 3 M sodium acetate, and 125 mM EDTA, and were sequenced with a Prism 3100 Genetic Analyser (ABI). Sequences were edited using SEQUENCHER v.4.1 (Genecodes). The type sequences were deposited in GenBank under DQ444270-DQ444275 accession numbers and EF183501-EF183502.

## EXTRACTION OF DNA FROM FRESH TISSUE

Total genomic DNA was extracted from fresh tissue of a *C. pulchristriata* that was field-collected in eastern Cambodia by B. L. Stuart (Table 1) using the Pure-Gene Animal Tissue DNA Isolation Protocol (Gentra Systems, Inc.). This sample represented the only fresh tissue of *Cyclemys* that had previously been extracted in the molecular laboratory where the historical DNA extractions and amplifications were performed. The extraction was amplified as a positive control in PCR reactions of historical DNA, and was

Table 1. Source	s of mitochondrial (mt) l	DNA of Cyclemys 1	turtles used in thi	is study		
Species	Museum voucher	Type status	Year collected	Condition	Locality	mtDNA sequence
atripons atripons bellii dentata oldhamii orbiculata ovata pulchristriata pulchristriata pulchristriata pulchristriata pulchristriata	USNM 81865 USNM 53423 USNM 53423 OUMNH 8513 BMNH 1946.1.22.62 BMNH 1947.3.5.63 BMNH 1947.3.4.26 OUMNH 8512 BMNH 1863.6.21.1 NMW 29525:4 NMW 29525:5 NMW 29525:5 SMF 7667 SMF 7667 SMF 7667 FMNH 259050	Holotype Paratype Holotype* Lectotype Paralectotype Syntype Holotype Paratype Paratype Paratype Paratype Paratype Paratype None	1929 1914 By 1834 $\dagger$ By 1834 $\dagger$ By 1856 $\$$ By 1856 $\$$ By 1862 $\P$ By 1863** By 1901 $\ddagger$ By 1901 $\ddagger$ By 1901 $\ddagger$ By 1901 $\ddagger$ By 1901 $\ddagger$ By 1901 $\ddagger$	Dried stuffed specimen Dried shell Dried shell Alcoholic specimen Dried shell Dried shell Dried shell Dried shell Alcoholic specimen Alcoholic specimen Alcoholic specimen Alcoholic specimen Alcoholic specimen Alcoholic specimen Alcoholic specimen Alcoholic specimen Alcoholic specimen Alcoholic specimen	Kao Kuap, Trat, Thailand Koh Chang, Trat, Thailand Unknown Java, Indonesia Mergui, Myanmar Lao Mountains, Thailand Unknown Sarawak, Malaysia†† Phuc-Son, Annam, Vietnam Phuc-Son, Annam, Vietnam Phuc-Son, Annam, Vietnam Annam, Vietnam Annam, Vietnam Annam, Vietnam	GenBank DQ444271 GenBank DQ444270 GenBank EF183501 GenBank DQ444272 GenBank DQ444273 GenBank DQ444273 GenBank EF183502 GenBank DQ444275 Failed to amplify Failed to amplify
Museum abbrev University Mus Smithsonian Ini *Das (2005) erry that it is the hc description of $C$ †This specimen $\ddagger$ The original B \$The original B \$The original B \$The original B \$The original B \$	iations: BMNH, The Nat atum of Natural History; titution; ZMH, Zoologisc oneously reported that O lotype of <i>C. bellii</i> . OUM <i>bellii.</i> is simultaneously a synt MNH catalogue number MNH catalogue number von the original spec year of the original spec lfred Russel Wallace. Ians Fruhstorfer, a Gern	ural History Muse SMF, Forschungs hes Museum Ham UMNH 8867 is th UMNH 8513 also matu ype of <i>C. orbiculai</i> 1828.5.12.1 implies 1862.8.18.20 implies 1862.8.18.20 implies ies description. an-Swiss natural	eum, London; FMI iinstitut und Natu nburg. te holotype of $C$ . $b$ ches the upper fig ches the upper fig sthat this specime ies that this specime ist who traveled t	NH, The Field Museum; NJ urmuseum Senckenberg, Fr <i>ellii</i> . The second author exu ure in plate 3, part 8 of Be so must have been collecte the nas been in the museur men has been in the museur men has been in the museur o Annam during a south-ee	<ul> <li>dW, Naturhistorisches Museum W ankfurt; USNM, National Museuu amined OUMNH 8513 and it bear ll (1832–1836), on which Gray (18 d by that year.</li> <li>d by that year.</li> <li>m since at least 1828.</li> <li>t since at least 1856.</li> <li>um since at least 1862.</li> <li>ust Asian expedition from 1899 to</li> </ul>	<ul> <li>ien; OUMNH, Oxford</li> <li>m of Natural History,</li> <li>rs an old label stating</li> <li>63) based his original</li> <li>63) based his original</li> <li>1901 (Lamas, 2005).</li> </ul>

**Table 2.** Oligonucleotide primers used to amplify and sequence part of the mitochondrial cytochrome b gene from *Cyclemys* turtles

Primer	Sequence
L-350cytb	5'-TTTGAGGTGCCACCGTCATC-3'
H-500cytb	5'-TCGTGGAGGAAGATTAAGTGTAC-3'
L-430cytb	5'-GGGATTCTCAGTAGACAATGC-3'
H-615cytb	5'-TTAGTAGGAGGGTGAGTATTAGGA-3'
L-560cytb	5'-CAGGATTAAACTCAAACGCTG-3'
H-730cytb	5'-GCATAGGCGAATAGGAAGTATC-3'

'L' and 'H' refer to light and heavy strands, respectively.

sequenced as described above. This sequence was deposited in GenBank under accession number DQ444276.

#### ALIGNMENT

In addition to the eight type specimens and one positive control tissue sequenced in this study, *Cyclemys* cyt *b* sequences were also obtained from three published sources. Thirty-nine sequences with a maximum length of 984 bp were obtained from Guicking *et al.* (2002), three sequences of 1140 bp (GenBank AY434577, AY434579, and AY434617) were obtained from Spinks *et al.* (2004), and one sequence of 1036 bp (GenBank AJ604513) was obtained from Schilde, Barth & Fritz (2004). Sequences from *Leucocephalon yuwonoi* (GenBank AY434608) and *Heosemys spinosa* (GenBank AY434578) were used as outgroups, based on the findings of Spinks *et al.* (2004). Sequences were aligned using SEQUENCHER v.4.1 (Genecodes).

#### PHYLOGENETIC ANALYSES

Phylogenies were reconstructed using maximumparsimony and maximum-likelihood optimality criteria, and mixed-model Bayesian inference.

Maximum-parsimony analysis was performed using PAUP\* 4.0b10 (Swofford, 2002). A heuristic search was performed with equal weighting of nucleotide substitutions, stepwise addition with 100 randomaddition replicates, and TBR branch swapping was limited to 1000 trees per replicate. Nodal support was evaluated with 500 nonparametric bootstrap pseudoreplications (Felsenstein, 1985) using the heuristic search option, with TBR branch swapping limited to 10 000 000 rearrangements per replicate.

Maximum-likelihood analysis was performed using GARLI 0.95 (Zwickl, 2006). Five independent analyses were performed using the default settings, the GTR + I + G model, and estimated base frequencies A = 0.31052, C = 0.30726, G = 0.11636, T = 0.26586,

rate matrix A-C = 1.04241, A-G = 8.18342, A-T = 0.4883, C-G = 0.00549, C-T = 12.0303, proportion of invariable sites = 0.0458648, and gammadistribution shape parameter = 0.283551. Nodal support was evaluated with 500 nonparametric bootstrap pseudoreplications.

Mixed-model Bayesian analysis was performed using MrBayes 3.1 (Ronquist & Huelsenbeck, 2003). The data were partitioned into first-codon, secondcodon, and third-codon positions. The model of sequence evolution that best described the three data partitions was inferred using the Akaike Information Criterion, as implemented in MODELTEST 3.7 (Posada & Crandall, 1998). The selected models were K81uf+G for the first-codon position partition. TrN + I for the second-codon position partition, and GTR+G for the third-codon position partition. The K81uf and TrN models are not implemented in MrBayes 3.1, and so the next more complex model available in the program (GTR) was used for those partitions. Four independent Bayesian analyses were performed. In each analysis, four chains were run for 10 000 000 generations using the default priors, trees were sampled every 2000 generations, and the first 25% of trees were discarded as 'burn-in'. A 50% majority rule consensus of the sampled trees was constructed to calculate the posterior probabilities of the tree nodes. Trace plots of clade probabilities were viewed using AWTY (Wilgenbusch, Warren & Swofford, 2004).

#### RESULTS

### HISTORICAL DNA SEQUENCES

All three overlapping cyt b fragments, consisting of a total of 376 bp, were obtained from the seven dried type specimens. Two overlapping cyt b fragments (from the primer pairs L-430cytb/H-615cytb and L-560cytb/H-730cytb), consisting of a total of 296 bp, were obtained from the alcoholic lectotype of *C. dentata*. None of the fragments were successfully amplified from the alcoholic type specimens of *C. pulchristriata* (Table 1).

The electropherograms of the obtained sequences had strong signals with very little or no background noise. Identical sequences were obtained in all overlapping fragments. The eight type sequences were genetically distinct from the positive control that represented the only fresh tissue of *Cyclemys* previously extracted in the same laboratory. Consequently, the sequences of the eight type specimens used in the analyses here appear to be authentic.

#### PHYLOGENETIC RELATIONSHIPS

The alignment contained no insertion-deletions and was unambiguous. Of the 1140 aligned characters,

297 were variable and 187 were parsimony informative. Strict consensus of the saved 100 000 equally most parsimonious trees (tree length, L = 435; consistency index, CI = 0.766; retention index, RI = 0.944) showed that these trees differed only by the arrangement of individuals on short-terminal branches. The score of the best-likelihood tree (-ln L 3565.994676) was within 0.043067 likelihood units of the best tree recovered in each of the other four runs, suggesting that the five runs had converged. The standard deviation of split frequencies among the four Bayesian runs was 0.004991, and trace plots of clade probabilities viewed using AWTY were relatively stationary. These two measures suggest that the four runs had converged, and that topologies were sampled in proportion to their true posterior probability distribution. The maximum-parsimony trees, maximum-likelihood tree, and Bayesian-consensus tree (Fig. 1) were highly consistent, and differed only by the arrangement of individuals on short-terminal branches.

Five major, well-supported clades were recovered (Clades A-E, Fig. 1). Clade A (Fig. 1) was sister to all other Cyclemys, and contained samples obtained from the Hong Kong pet trade (Guicking et al., 2002). These were reported to have originated from Kachin, Myanmar, and were identified as an undescribed species morphologically resembling C. oldhamii (Guicking et al., 2002). None of the type sequences that we obtained belong to this clade. Clade B (Fig. 1) contained samples identified as two subspecies of C. shanensis by Guicking et al. (2002), as C. shanensis by Schilde et al. (2004), as C. dentata and C. tcheponensis by Spinks et al. (2004), and the lectotype and paralectotype of C. oldhamii. Clade C (Fig. 1) contained samples identified as C. atripons and C. pulchristriata by Guicking et al. (2002), and the positive control sample of C. pulchristriata from eastern Cambodia. Clade D (Fig. 1) contained samples identified as C. dentata and C. oldhamii by Guicking et al. (2002), the lectotype of *C. dentata*, the syntype of C. orbiculata, the holotype of C. bellii, and the holotype of C. ovata. Clade E (Fig. 1) contained samples identified by Guicking et al. (2002) as an undescribed species morphologically resembling C. atripons and C. pulchristriata, and the holotype and paratype of C. atripons. Species with yellow plastra (Clades C-E, Fig. 1) form a well-supported monophyletic group, but species with dark plastra (Clades A-B, Fig. 1) are paraphyletic (Fig. 1).

# DISCUSSION

Our phylogenetic analysis of mitochondrial DNA sequences obtained from museum type specimens of *Cyclemys* greatly aids the resolution of species boundaries in the genus. Specifically, our phylogenetic

hypothesis supports the recognition of four named species (C. atripons, C. dentata, C. oldhamii, and C. pulchristriata), and a fifth species of unknown geographical origin. The substantial differences in identifications among Guicking et al. (2002), Spinks et al. (2004), and this study incorporating type sequences, support the notion that Cyclemys of unknown provenance are not reliably identified to species solely on the basis of morphology. The corrected taxonomy based on the type sequences (Fig. 1) shows that many samples were misidentified in previous studies (Guicking et al., 2002; Spinks et al., 2004). Unfortunately, the lack of voucher specimens associated with some (Guicking et al., 2002) or all (Spinks et al., 2004) of the genetic material analysed in those studies prevents re-evaluating identifications in the light of our new findings with type sequences.

The absence of substantial genetic variation among the types of C. bellii, C. dentata, C. orbiculata, and C. ovata corroborates the hypothesis based on morphology (Fritz et al., 1997) that C. bellii, C. orbiculata, and C. ovata are junior synonyms of C. dentata. the oldest available name. Likewise, the absence of substantial genetic variation among the types of C. oldhamii and samples identified as C. shanensis (including topotypes of C. s. shanensis from Lake Inlé, Myanmar) and C. tcheponensis suggests that C. shanensis and C. tcheponensis are junior synonyms of C. oldhamii, the oldest available name. Cyclemys oldhamii, C. shanensis, and C. tcheponensis differ morphologically only by the retention or loss of head stripes with maturity (Fritz et al., 1997; Guicking et al., 2002), and adopting this taxonomy implies that the retention or loss of head stripes with maturity represents only geographical variation within a species.

Cyclemys atripons and C. pulchristriata are morphologically indistinguishable, but are paraphyletic with respect to C. dentata. Because we were unsuccessful in obtaining DNA from the type series of C. pulchristriata (Table 1), our identification of the C. pulchristriata clade is tentatively based on the assumption that the positive control sample belongs to this species. This sample was field-collected in hilly eastern Cambodia by B. L. Stuart, in the same Annamite (Truong Son) Mountain range that contains the type locality of C. pulchristriata in central Vietnam. Stuart & Platt (2004) tentatively referred this specimen to C. atripons, on the basis that most of the Cambodian Cyclemys specimens with yellow plastra they examined originated from the mountains of south-western Cambodia, close to the type locality of C. atripons, and that no morphological differences among the specimens were apparent. Stuart, Sok & Neang (2006b) identified additional specimens from eastern Cambodia as C. atripons to maintain consis-



**Figure 1.** Fifty-percent majority-rule consensus phylogram resulting from mixed-model Bayesian analysis of the mitochondrial cytochrome *b* gene from Asian leaf turtles (*Cyclemys*). Maximum-likelihood and maximum-parsimony analysis recovered the same topology, and differed only by the arrangement of individuals on short-terminal branches. Numbers next to nodes are Bayesian posterior probabilities/maximum-likelihood bootstrap values/parsimony bootstrap values. Haplotypes in boldface were sequenced in this study, haplotypes marked with an asterisk (\*) were obtained from Spinks *et al.* (2004), the haplotype marked with two asterisks (\*\*) was obtained from Schilde *et al.* (2004), and the remaining haplotypes were obtained from Guicking *et al.* (2002). Species identifications of haplotypes obtained from Guicking *et al.* (2002), Schilde *et al.* (2004), and Spinks *et al.* (2004) are enclosed in single quotations. Our recommended taxonomy is on the right.

tency with Stuart & Platt (2004). Sequences from type material of *C. pulchristriata* or from fresh topotypic samples are needed to identify the *C. pulchristriata* lineage with certainty. It is clear, however, from our results that the lineage interpreted by Guicking et al. (2002) to be a cryptic, undescribed species morphologically resembling *C. atripons* and *C. pulchristriata* ('Cyclemys n. sp. 2') is *C. atripons*.

The lineage interpreted by Guicking *et al.* (2002) to be a cryptic, undescribed species morphologically

resembling C. oldhamii ('Cyclemys n. sp. 1') does not match any of the type sequences obtained in this study, and may indeed represent an undescribed species. We did not attempt to sequence the type material of three described taxa of Cyclemys that are currently considered to be junior synonyms of other species, and which exhibit a dark plastron like that of 'Cyclemys n. sp. 1': C. tcheponensis, C. dhor shanensis, and C. tiannanensis. The sequences obtained from Guicking et al. (2002) include samples from Lake Inlé, Myanmar, the type locality of C. dhor shanensis (referred to as C. s. shanensis in Guicking et al., 2002) and from Lao Bao, Vietnam, near to the type locality of C. tcheponensis (referred to as C. s. tcheponensis in Guicking et al., 2002), and so it is unlikely that 'Cyclemys n. sp. 1' represents either of these two taxa. Cyclemys tiannanensis has not yet been characterized molecularly, and its relationship to 'Cyclemys n. sp. 1' remains unevaluated. Unfortunately, the provenance of 'Cyclemys n. sp. 1' is unknown. These samples were obtained from a Hong Kong pet dealer who reported that they originated from Kachin Province. Myanmar (Guicking et al., 2002). During the past two decades, many newly described turtle species were based on specimens purchased from a Hong Kong pet dealer (Parham et al., 2001; Dalton, 2003). Some of these newly described species have been shown to be hybrids of other, better known species (Parham et al., 2001; Wink, Guicking & Fritz, 2001; Spinks et al., 2004; Stuart & Parham, 2007), whereas others have been shown to represent distinct, evolutionary lineages that remain unknown in the wild (Parham et al., 2004). Because the mitochondrial DNA of 'Cyclemys n. sp. 1' does not match any other known taxon (except potentially C. tiannanensis), this distinct lineage is unlikely to be another example of a pet-trade species with a recent hybrid origin. In many cases, the provenance supplied by the Hong Kong pet dealer for specimens have been questioned (de Bruin & Artner, 1999; Parham & Li, 1999; Dalton, 2003; Shi et al., 2005), underscoring the urgent need for scientific fieldwork.

Although we hypothesize that the major mitochondrial DNA lineages represent species of *Cyclemys*, one apparent example in our dataset of introgression between mitochondrial DNA lineages suggests that these lineages may be imperfect proxies for species. Mitochondrial DNA sequences obtained by Guicking *et al.* (2002) from two turtles with black plastra purchased in local trade at Penang, Peninsular Malaysia, and identified by those authors based on morphology as *C. oldhamii*, are deeply nested within a clade of turtles with yellow plastra referred here to *C. dentata* (Fig. 1). A nuclear DNA dataset (ISSR-PCR) presented by Guicking *et al.* (2002) on the same samples readily distinguished the Penang turtles from *C.*  *dentata*, supporting the hypothesis that the Penang samples are an example of mitochondrial DNA introgression. Cyclemys oldhamii and C. dentata occur in sympatry in Peninsular Malaysia (Fritz et al., 1997), but owing to the trade origin of the Penang samples, the introgression may alternatively have occurred in captivity. Field sampling in Peninsular Malaysia should offer insight into this topic. A nuclear DNA dataset of the type sequences would also provide an independent estimate of relationships and species boundaries in Cyclemys. However, obtaining nuclear DNA sequences from degraded samples is more challenging than obtaining mitochondrial DNA sequences, probably because mitochondrial DNA occurs in higher copy number in the cell, and is therefore more likely to be retrieved (Hofreiter et al., 2001; Huynen et al., 2003; Isenberg, 2005).

Sampling genetic material from a museum specimen necessarily requires removing part of the specimen, and there is justifiably great concern that damage to the specimen be kept to a mimimum, particularly with those specimens that are as irreplaceable as types (Graves & Braun, 1992; Mundy, Unitt & Woodruff, 1997; Rohland, Siedel & Hofreiter, 2004; Wisely et al., 2004; Asher & Hofreiter, 2006). New protocols for obtaining DNA from museum specimens are commendably seeking to minimize such damage (Mundy et al., 1997; Rohland et al., 2004; Wisely et al., 2004; Asher & Hofreiter, 2006), but still modify specimens in some manner. We submit that sampling genetic material from freshly collected specimens of turtles also requires destructive sampling by removing tissue (typically muscle or liver) from the carcass. As with freshly collected specimens, in this study we aimed to sample tissues that were subjectively deemed to have limited phylogenetic utility. These included small pieces of skin, muscle, bone, or connective tissue, usually in the form of unrecognizable organic material adhering to the inside of dried specimens. The most drastic sampling (and ironically the least successful) entailed removing a femur bone from a leg that was retracted within the shell of some alcoholic specimens. Efforts should continue to find new means to extract DNA with minimal damage to specimens, and curators should continue to critically review requests to destructively sample specimens in their care (Graves & Braun, 1992). However, our ability to sample type specimens of Cyclemys in this manner enabled us to reveal new phylogenetic characters from critically important specimens, including one notably collected by Alfred Russel Wallace.

The type sequences generated in this study provide useful reference points for future phylogenetic studies of *Cyclemys* because their species identifications are unambiguous. New efforts to resolve *Cyclemys* species boundaries and relationships should be directed towards obtaining mitochondrial DNA sequences from *C. pulchristriata* and *C. tiannanensis* type material, obtaining nuclear DNA sequences from all type material, and especially, obtaining field-collected, vouchered, genetic material of known provenance. The genus *Cyclemys* contains multiple, distinct evolutionary lineages that each warrant recognition and protection, but an effective conservation strategy requires knowing where these lineages exist in nature.

# ACKNOWLEDGEMENTS

The following people (and institutions) made this study possible by sampling and loaning tissue from type specimens in their care: George Zug (National Museum of Natural History, Smithsonian Institution), Colin McCarthy (The Natural History Museum, London), Richard Gemel and Franz Tiedemann (Naturhistorisches Museum Wien), Gunther Köhler (Forschungsinstitut und Naturmuseum Senckenberg, Frankfurt), Tom Kemp and Malgosia Nowak-Kemp (Oxford University Museum of Natural History), and Jakob Hallermann (Zoologisches Museum Hamburg). The specimen from Mondolkiri, Cambodia, was collected under the auspices of the Wildlife Conservation Society/Ministry of Agriculture, Forestry and Fisheries/Ministry of Environment Collaborative Program with field assistance from An Dara and Suon Phalla. Editha Schubert and Holger Dathe (Deutsches Entomologisches Institut, Müncheberg) and Hannes Baur (Naturhistorisches Museum Bern) provided biographical information on Hans Fruhstorfer. Sequencing was performed in The Field Museum's Pritzker Laboratory for Molecular Systematics and Evolution operated with support from the Pritzker Foundation. Rob Vogelbacher and the DePaul University Bioinformatics Group, in conjunction with the Illinois Bio-Grid, executed the Bayesian analysis on a computer cluster. The work was supported by a grant from The John D. and Catherine T. MacArthur Foundation to the Field Museum. James Parham (California Academy of Sciences) and two anonymous reviewers improved the manuscript.

#### REFERENCES

Amato G, Egan MG, Schaller GB, Baker RH, Rosenbaum HC, Robichaud WG, DeSalle R. 1999. Rediscovery of Roosevelt's barking deer (Muntiacus rooseveltorum). Journal of Mammalogy 80: 639–643.

Annandale N. 1918. Chelonia and Batrachia of the Inlé Lake. Records of the Indian Museum 14: 67–69.

Artner H. 1998. Nomenklatur aktuell. Emys 5: 33.

- Asher RJ, Hofreiter M. 2006. Tenrec phylogeny and the noninvasive extraction of nuclear DNA. Systematic Biology 55: 181–194.
- Austin JJ, Arnold EN. 2001. Ancient mitochondrial DNA and morphology elucidate an extinct island radiation of Indian Ocean giant tortoises (*Cylindraspis*). Proceedings of the Royal Society B 268: 2515–2523.
- Austin JJ, Arnold EN, Bour R. 2002. The provenance of type specimens of extinct Mascarene Island giant tortoises (*Cylindraspis*) revealed by ancient mitochondrial DNA sequences. Journal of Herpetology **36**: 280–285.
- Austin JJ, Arnold EN, Bour R. 2003. Was there a second adaptive radiation of giant tortoises in the Indian Ocean? Using mitochondrial DNA to investigate speciation and biogeography of *Aldabrachelys* (Reptilia, Testudinidae). *Molecular Ecology* 12: 1415–1424.
- Austin JJ, Arnold EN, Jones CG. 2004. Reconstructing an island radiation using ancient and recent DNA: the extinct and living day geckos (*Phelsuma*) of the Mascarene islands. *Molecular Phylogenetics and Evolution* **31:** 109–122.
- Austin JJ, Melville J. 2006. Incorporating historical museum specimens into molecular systematic and conservation genetics research. *Molecular Ecology Notes* 6: 1089– 1092.
- Bell T. 1832–1836. A monograph of the Testudinata. London: Samuel Highley.
- Bell T. 1834. Characters of a new genus of freshwater tortoise (*Cyclemys*). Proceedings of the Zoological Society of London 1834: 17.
- **Bourret R. 1939.** Notes herpétologiques sur l'Indochine française. XVI. Tortues de la collection du Laboratoire des Sciences naturelles de l'Université. Description d'une espèce nouvelle. Annexe au Bulletin Général de l'Instruction Publique **1938**: 5–12.
- de Bruin RWF, Artner HG. 1999. On the turtles of Hainan Island, southern China. *Chelonian Conservation and Biology* 3: 479–486.
- Cooper A, Poinar HN. 2000. Ancient DNA: do it right or not at all. Science 289: 1139.
- Dalton R. 2003. Mock turtles. Nature 423: 219-220.
- **Das I. 2005.** Asian turtles in the collection of the Oxford University Museum, including a list of type specimens. *Herpinstance* **3:** 4–7.
- van Dijk PP, Stuart BL, Rhodin AGJ, eds. 2000. Asian turtle trade: proceedings of a workshop on conservation and trade of freshwater turtles and tortoises in Asia. Chelonian Research Monographs 2. Lunenburg: Chelonian Research Foundation.
- Felsenstein J. 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* **39**: 783–791.
- Fleischer RC, Kirchman JJ, Dumbacher JP, Bevier L, Dove C, Rotzel NC, Edwards SV, Lammertink M, Miglia KJ, Moore WS. 2006. Mid-Pleistocene divergence of Cuban and North American ivory-billed woodpeckers. *Biology Letters* 2: 466–469.
- Fritz U, Gaulke M, Lehr E. 1997. Revision der südostasiatischen Dornschildkröten-Gattung *Cyclemys* Bell, 1834, mit Beschreibung einer neuen Art. *Salamandra* 33: 183–212.

- Fritz U, Guicking D, Wink M, Lehr E. 2001. Sind Cyclemys atripons Iverson & McCord, 1997 und Cyclemys pulchristriata Fritz, Gaulke & Lehr, 1997 identisch? Sauria 23: 33–38.
- Fritz U, Obst FJ. 1999. Neue Schildkröten aus Südostasien Teil II. Bataguridae (Cyclemys, Heosemys, Mauremys, Ocadia, Pyxidea, Sacalia) und Trionychidae. Sauria 21: 11–26.
- Fritz U, Ziegler T. 1999. Contribution to the knowledge of Cyclemys tcheponensis (Bourret, 1939) and the distribution of Cyclemys in the Indochinese region (Reptilia: Testudines: Bataguridae). Revue français d'Aquariologie 26: 71–78.
- Fuchs J, Fjeldså J, Pasquet E. 2005. The use of mitochondrial and nuclear sequence data in assessing the taxonomic status of the endangered Uluguru Bush Shrike *Malaconotus alius*. *Ibis* 147: 717–724.
- Graves GR, Braun MJ. 1992. Museums: storehouses of DNA? Science 255: 1335–1336.
- Gray JE. 1831. Synopsis Reptilium or short descriptions of the species of reptiles. Part I. Cataphracta, tortoises, crocodiles, and enaliosaurians. London: Treuttel, Wurtz, and Co.
- Gray JE. 1863. Observations on the box tortoises, with the descriptions of three new Asiatic species. *Proceedings of the Zoological Society of London* 1863: 173–179.
- Guicking D, Fritz U, Wink M, Lehr E. 2002. New data on the diversity of the Southeast Asian leaf turtle genus Cyclemys Bell, 1834. Molecular results (Reptilia: Testudines: Geoemydidae). Faunistiche Abhandlungen Staatliches Museum für Tierkunde Dresden 23: 75–86.
- Handt O, Höss M, Krings M, Pääbo S. 1994. Ancient DNA: methodological challenges. *Experientia* 50: 524–529.
- Hassanin A, Ropiquet A. 2004. Molecular phylogeny of the tribe Bovini (Bovidae, Bovinae) and the taxonomic status of the Kouprey, Bos sauveli Urbain 1937. Molecular Phylogenetics and Evolution 33: 896–907.
- Hofreiter M, Serre D, Poinar HN, Kuch M, Pääbo S. 2001. Ancient DNA. Nature Reviews. Genetics 2: 353–359.
- Hughey JR, Silva PC, Hommersand MH. 2001. Solving taxonomic and nomenclatural problems in Pacific Gigartinaceae (Rhodophyta) using DNA from type material. *Journal of Phycology* 37: 1091–1109.
- Huynen L, Millar CD, Scofield RP, Lambert DM. 2003. Nuclear DNA sequences detect species limits in ancient DNA. *Nature* 425: 175–178.
- Isenberg AR. 2005. Forensic mitochondrial DNA analysis. In: Saferstein R, ed. *Forensic science handbook*, Vol. II, 2nd edn. Upper Saddle River, NJ: Pearson Education, Inc., 297– 327.

IUCN. 2004. Red list of threatened species. Available at http:// www.redlist.org

IUCN/SSC Tortoise and Freshwater Turtle Specialist Group and Asian Turtle Trade Working Group. 2000. Recommended changes to 1996 IUCN Red List status of Asian turtle species. In: van Dijk PP, Stuart BL, Rhodin AGJ, eds. Asian turtle trade: proceedings of a workshop on conservation and trade of freshwater turtles and tortoises in Asia. Chelonian Research Monographs 2. Lunenburg: Chelonian Research Foundation, 156–164.

- **Iverson JB, McCord WP. 1997.** A new species of *Cyclemys* (Testudines: Bataguridae) from Southeast Asia. *Proceedings* of the Biological Society of Washington **110:** 629–639.
- Johnson JA, Watson RT, Mindell DP. 2005. Prioritizing species conservation: does the Cape Verde kite exist? *Proceedings of the Royal Society B* 272: 1365–1371.
- Kearney M, Stuart BL. 2004. Repeated evolution of limblessness and digging heads in worm lizards revealed by DNA from old bones. *Proceedings of the Royal Society B* 271: 1677–1683.
- Kohlsdorf T, Wagner GP. 2006. Evidence for the reversibility of digit loss: a phylogenetic study of limb evolution in *Bachia* (Gymnophthalmidae: Squamata). *Evolution* 60: 1896–1912.
- Kou Z. 1989. Cyclemys from Yunnan, a description of a new species and a new record for China (Testudinata: Emydidae). In: Matsui M, Hikida T, Goris RC, eds. Current herpetology in East Asia. Kyoto: The Herpetological Society of Japan, 193–197.
- Krings M, Stone A, Schmitz RW, Krainitzki H, Stoneking M, Pääbo S. 1997. Neandertal DNA sequences and the origin of modern humans. *Cell* 90: 19–30.
- Lamas G. 2005. A bibliography of the zoological publications of Hans Fruhstorfer (1866\*–1922†). Entomofauna 26: 57–100.
- Maddison WP. 1997. Gene trees in species trees. Systematic Biology 46: 523–536.
- Moore WS. 1995. Inferring phylogenies from mtDNA variation: mitochondrial-gene tree versus nuclear-gene trees. *Evolution* 49: 718–726.
- Mundy NI, Unitt P, Woodruff DS. 1997. Skin from feet of museum specimens as a non-destructive source of DNA for avian genotyping. *The Auk* 114: 126–129.
- Olson LE, Hassanin A. 2003. Contamination and chimerism are perpetuating the legend of the snake-eating cow with twisted horns (*Pseudonovibos spiralis*). A case study of the pitfalls of ancient DNA. *Molecular Phylogenetics and Evolution* 27: 545–548.
- Parham JF, Li D. 1999. A new locality for Cuora pani Song 1984 with comments on its known range. Asiatic Herpetological Research 8: 111–113.
- Parham JF, Simison WB, Kozak KH, Feldman CR, Shi H. 2001. New Chinese turtles: endangered or invalid? A reassessment of two species using mitochondrial DNA, allozyme electrophoresis and known locality specimens. *Animal Conservation* 4: 357–367.
- Parham JF, Stuart BL, Bour R, Fritz U. 2004. Evolutionary distinctiveness of the extinct Yunnan box turtle (*Cuora yunnanensis*) revealed by DNA from an old museum specimen. Proceedings of the Royal Society B (Biology Letters Supplement) 271: S391–S394.
- Payne RB, Sorenson MD. 2002. Museum collections as sources of genetic data. Bonner Zoologische Beiträge 51: 97–104.
- Posada D, Crandall KA. 1998. Modeltest: testing the model of DNA substitution. *Bioinformatics* 14: 817–818.
- Rohland N, Siedel H, Hofreiter M. 2004. Nondestructive DNA extraction method for mitochondrial DNA analyses of museum specimens. *Biotechniques* 36: 814–821.

- Ronquist F, Huelsenbeck JP. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572–1574.
- Roy MS, Girman DJ, Taylor AC, Wayne RK. 1994. The use of museum specimens to reconstruct the genetic variability and relationships of extinct populations. *Experientia* 50: 551–557.
- Schilde M, Barth D, Fritz U. 2004. An Ocadia sinensis x Cyclemys shanensis hybrid (Testudines: Geoemydidae). Asiatic Herpetological Research 10: 120–125.
- Shi H, Parham JF, Simison WB, Wang J, Gong S, Fu B. 2005. A report on the hybridization between two species of threatened Asian box turtles (Testudines: *Cuora*) in the wild on Hainan Island (China) with comments on the origin of '*serrata*'-like turtles. *Amphibia-Reptilia* 26: 377–381.
- Sorenson MD, Cooper A, Paxinos EE, Quinn TW, James HF, Olson SL, Fleischer RC. 1999. Relationships of the extinct moa-nalos, flightless Hawaiian waterfowl, based on ancient DNA. *Proceedings of the Royal Society B* 266: 2187– 2193.
- Spinks PQ, Shaffer HB, Iverson JB, McCord WP. 2004. Phylogenetic hypotheses for the turtle family Geoemydidae. *Molecular Phylogenetics and Evolution* 32: 164–182.
- Stuart BL, Inger RF, Voris HK. 2006a. High level of cryptic species diversity revealed by sympatric lineages of Southeast Asian forest frogs. *Biology Letters* 2: 470–474.
- Stuart BL, Parham JF. 2004. Molecular phylogeny of the critically endangered Indochinese box turtle (*Cuora galbinifrons*). Molecular Phylogenetics and Evolution 31: 164– 177.
- Stuart BL, Parham JF. 2007. Recent hybrid origin of three rare Chinese turtles. *Conservation Genetics* 8: 169– 175.
- Stuart BL, Platt SG. 2004. Recent records of turtles and tortoises from Laos, Cambodia, and Vietnam. Asiatic Herpetological Research 10: 129–150.
- Stuart BL, Sok K, Neang T. 2006b. A collection of

amphibians and reptiles from hilly eastern Cambodia. Raffles Bulletin of Zoology 54: 129–155.

- Stuart BL, Timmins RJ, Hendrie DB, Lieng S, Chun S, Hout P, Heng K, Touch TS, Prak HL, Chul T, Compton J, Holloway R. 2000. Turtle trade in Indochina: regional summary (Cambodia, Laos, and Vietnam). In: van Dijk PP, Stuart BL, Rhodin AGJ, eds. Asian turtle trade: proceedings of a workshop on conservation and trade of freshwater turtles and tortoises in Asia. Chelonian Research Monographs 2. Lunenburg: Chelonian Research Foundation, 74–76.
- **Swofford DL. 2002.** *PAUP\*: phylogenetic analysis using parsimony \* (and other methods)*, Version 4.0b10. Sunderland: Sinauer Associates.
- Tautz D, Arctander P, Minelli A, Thomas RH, Vogler AP. 2003. A plea for DNA taxonomy. Trends in Ecology and Evolution 18: 70–74.
- Wayne RK, Leonard JA, Cooper A. 1999. Full of sound and fury: the recent history of ancient DNA. Annual Review of Ecology and Systematics 30: 457–477.
- Wilgenbusch JC, Warren DL, Swofford DL. 2004. AWTY: a system for graphical exploration of MCMC convergence in Bayesian phylogenetic inference. Available at http:// ceb.csit.fsu.edu/awty
- Wink M, Guicking D, Fritz U. 2001. Molecular evidence for hybrid origin of *Mauremys iversoni* Pritchard et McCord, 1991, and *Mauremys pritchardi* McCord, 1997 (Reptilia: Testudines: Bataguridae). Zoologische Abhandlungen Staatliches Museum für Tierkunde Dresden 51: 41-49.
- Wisely SM, Maldonado JE, Fleischer RC. 2004. A technique for sampling ancient DNA that minimizes damage to museum specimens. *Conservation Genetics* 5: 105–107.
- Zwickl DJ. 2006. Genetic algorithm approaches for the phylogenetic analysis of large biological sequence datasets under the maximum likelihood criterion. PhD Dissertation, The University of Texas at Austin, USA. [Genetic Algorithm for Rapid Likelihood Inference software available at http:// www.bio.utexas.edu/faculty/antisense/garli/Garli.html].